

FOR ANIMAL TREATMENT ONLY
RESTRICTED VETERINARY MEDICINE



TUBERCULIN 30,000 IU per mL

Registered to: AsureQuality Limited,
Level 1, 7a Pacific Rise, Mt Wellington 1060,
New Zealand
Tel: +64 9 573 8000
Fax: +64 9 573 8118
Registered pursuant to ACVM Act 1997
No. A10768.

See www.foodsafety.govt.nz for
registration conditions



STORE AT 2 TO 8 DEGREES CELSIUS
DO NOT FREEZE
PROTECT FROM LIGHT
KEEP OUT OF REACH OF CHILDREN

W4599

PB3000PAU-01-MAY2021

FOR ANIMAL TREATMENT ONLY
RESTRICTED VETERINARY MEDICINE



TUBERCULIN 30,000 IU per mL

Active Ingredients: Purified Protein Derivative of
Mycobacterium bovis, strain AN5, 30,000 IU/mL.

Indications: For use as an aid in the diagnosis of
mycobacterial infection in cattle.

Directions for Use: For use under the authority or
prescription of a veterinary surgeon or by persons
approved under the Australian Biosecurity Act 2015
and Australian Export Control Act 1982 when
applying cattle tuberculin tests.

Test Application

The following instructions are guidelines only:
Injections of 0.1 mL must be made intradermally.
It is recommended syringes are calibrated to 0.1
mL and the needle of the syringe is 22-26 gauge
and 3-4 mm long. Syringes should be kept clean
and sterilised. Contamination of the syringe with
disinfectants or alcohol may interfere with testing.
When filling syringes, care must be taken to prevent
contamination of vials.

Preferred site of injection:

Cattle: caudal fold for the single intradermal test
and the cervical site for the comparative cervical
test (CCT).

When using the cervical sites, care is required
to evenly clip hair close to the skin surface
(2mm or less mean length) prior to injecting.
The recommended size of the clipped area is
10 x 10 cm for each injection site.

When applying a CCT use Observe™ Bovine
Tuberculin 30,000 IU per mL in conjunction with
Observe™ Avian Tuberculin 25,000 IU per mL.

Test Interpretation:

It is recommended the test be read 72 hours
after injection.

Skin test reactions are to be classified in accordance
with the requirements of OSPRI, the TBfree New
Zealand programme and the National Operational
Plan pursuant to the Biosecurity Act 1993 and the
Biosecurity (National Bovine Tuberculosis Pest
Management Plan) Order 1998 (the Order).

Retesting: To avoid desensitisation, a retest using
bovine and/or avian PPD tuberculin should not be
applied until 60 days after any previous tuberculin
test.

Product Information

Withholding period: Nil

Storage: Refrigerate (2°C to 8°C). Do not freeze.
Protect from light - keep vials in closed box.

Transportation: May be transported at
2°C to 37°C for a period not longer than 14 days.
Do not freeze.

Use: Do not use after the expiry date. Use
immediately after opening. Discard unused
contents.

Method of Disposal: Unused, part-used and
empty vials should be destroyed according to local
regulations.

Safety Precautions: In the case of accidental
self-injection, an area of intense local irritation may
develop for up to 96 hours later.

If irritation occurs, you are advised to consult your
medical practitioner.

Store out of the reach of children.

Manufactured by: Prionics Lelystad B.V.,
Platinastraat 33, 8211 AR Lelystad,
The Netherlands

Registered to: AsureQuality Limited,
Level 1, 7a Pacific Rise, Mt Wellington 1060,
New Zealand

Tel: +64 9 573 8000
Fax: +64 9 573 8118

Registered pursuant to ACVM Act 1997
No. A10768. See www.foodsafety.govt.nz
for registration conditions.

PB3000PAU-01-MAY2021